

Phase transitions and exchange interactions in the $\text{SmCr}_3(\text{BO}_3)_4$ crystal

Dobretsova E., Boldyrev K., Popova M., Gavrilkin S., Mukhin A., Ivanov V., Mal'tsev V., Leonyuk N., Malkin B.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© The Authors, published by EDP Sciences. Spectroscopic investigation and specific heat and magnetic susceptibility measurements of $\text{SmCr}_3(\text{BO}_3)_4$ crystals were performed. The spectra of the Sm^{3+} and Cr^{3+} ions in samarium chromium borate were calculated and parameters of the exchange interactions between the nearest chromium ions, chromium and samarium ions were determined. Three phase transitions were observed at the temperatures $T_1 = 7.8 \pm 0.5$ K, $T_2 = 6.7 \pm 0.5$ K, and $T_3 = 4.3 \pm 0.2$ K, their nature is discussed. The crystal structures with different space symmetry groups $R\bar{3}2$ and $C2/c$ were found to coexist in $\text{SmCr}_3(\text{BO}_3)_4$ single crystal.

<http://dx.doi.org/10.1051/epjconf/201713202008>

References

- [1] E.A. Popova et al., Phys. Rev. B 75, 054446 (2007).
- [2] L. Gondek et al., J. Sol. St. Chem. 210, 30 (2014).
- [3] K.N. Boldyrev, E.P. Chukalina, N.I. Leonyuk, Phys. Sol. St. 50, 1681 (2008).
- [4] M.N. Popova et al., Phys. Rev. B 75, 224435 (2007).